

## Activity-dependent Outgrowth of Neurons and Overshoot Phenomena in Developing Neural Networks

A. VAN OUYEN AND J. VAN PELT

*Netherlands Institute for Brain Research, Meibergdreef 33, 1105 AZ Amsterdam, The Netherlands*

(Received on 22 January 1993, Accepted in revised form on 17 May 1993)

During the development of the nervous system, all kinds of structural elements such as neurons, neuritic extensions and synapses are initially overproduced (so-called overshoot phenomena). Neurite outgrowth has been found to be regulated by electrical activity of the neuron. High levels of activity, resulting in high intracellular calcium concentrations, cause neurites to retract whereas low levels of activity, and consequently low calcium concentrations, allow further outgrowth. In this article, we have studied the implications of such activity-dependent neurite outgrowth for network formation, using a distributed simulation model. The model consists of initially disconnected cells that organize themselves into a network under the influence of their intrinsic activity. A neuron is modelled as a neuritic field, the growth of which depends upon its own level of activity, and the neurons become connected when their fields overlap. It is demonstrated here that activity-dependent outgrowth in combination with a neuronal response function with some form of firing threshold—which gives rise to a hysteresis effect—is sufficient to cause an overshoot with respect to connectivity or number of synapses. As a consequence of such hysteresis, the network connectivity at which a phase transition occurs from the quiescent to the activated state must be higher than that for maintaining activity at a level where neuritic fields and connectivity remain constant. A developing network will therefore first increase its connectivity until it becomes activated, upon which the neurites begin to retract. Connectivity then decreases until the equilibrium value is reached, thus causing the growth curve to exhibit overshoot. The results are robust under various alternative formulations of the model, and show certain similarities with findings in developing cultures of dissociated nerve cells, namely a transient overproduction of synapses and the existence of a transition period in which increasing electrical activity is associated with retraction of neurites.

### 1. Introduction

Neurons become assembled into functional neural networks during development. Among the many factors influencing the ultimate structure and function of the nervous system, electrical activity plays a pivotal role (for review see Fields & Nelson, 1992). Many mechanisms that determine neuronal connectivity such as neurite outgrowth, growth cone behaviour, naturally occurring cell death, trophic factors, synaptogenesis, elimination of synapses, and changes in synaptic strength have been found to be modulated by electrical activity.

### 1.1. NEURITE OUTGROWTH

Electrical activity of the neuron reversibly arrests neurite outgrowth (or produces retraction) and changes growth cone morphology (Cohan & Kater, 1986; Fields *et al.*, 1990a; Schilling *et al.*, 1991). Similarly, depolarizing media and neurotransmitters affect neurite outgrowth of many cell types (e.g. Sussdorf & Campenot, 1986; Lankdorf *et al.*, 1987; McCobb *et al.*, 1988; Lipton & Kater, 1989; Mattson & Kater, 1989; Todd, 1992), with excitatory neurotransmitters inhibiting outgrowth and inhibitory ones stimulating outgrowth. Electrical activity and

neurotransmitters probably regulate neurite outgrowth by affecting the calcium concentration in growth cones (Cohan *et al.*, 1987; Fields *et al.*, 1990b; Kater & Mills, 1991). This has led to the calcium theory of neurite outgrowth (e.g. Kater *et al.*, 1988, 1990; Kater & Guthrie, 1990), which posits that low intracellular calcium concentrations ( $[Ca^{2+}]_i$ ) stimulate outgrowth, higher concentrations cause a cessation of outgrowth, and still higher concentrations lead to regression of neurites. Thus, all factors that change  $[Ca^{2+}]_i$  (such as action potentials or other forms of depolarization and neurotransmitter actions) are potentially able to affect neurite outgrowth. Alterations in  $[Ca^{2+}]_i$  have also been implicated in the development of dendritic morphology (Kater *et al.*, 1990). Because of the morphological changes that accompany changes in  $[Ca^{2+}]_i$ , and the large number of signals that influence it, mechanisms regulating  $[Ca^{2+}]_i$  have been proposed to represent a major means by which entire patterns of neuronal circuitry can be specified (Lipton & Kater, 1989).

Applied electric fields, too, influence nerve growth, with respect both to branching (McCaig, 1990a) and to the rate of elongation (McCaig, 1990b).

### 1.2. CELL DEATH

Neuronal activity may play an important role in the regulation of naturally occurring cell death (e.g. Brenneman & Eiden, 1986; Lipton, 1986; Oppenheim, 1991; Ferrer *et al.*, 1992). The neurotrophic theory, which asserts that neurons depend upon, and compete for, neurotrophic factors that are released only in limited amounts by their targets (Hamburger & Oppenheim, 1982; Purves, 1988), provides a framework for understanding cell death during normal ontogeny. The synthesis of the neurotrophic factors NGF (nerve growth factor) and BDNF (brain derived neurotrophic factor) by neurons in the central nervous system has been found to be regulated by neuronal activity (Zafra *et al.*, 1990; Lu *et al.*, 1991; Thoenen, 1991). The degree of neuronal dependence on NFG may also be modulated by electrical activity, since it depends on  $[Ca^{2+}]_i$  (Koike & Tanaka, 1991).

### 1.3. SYNAPSES

The refinement of imprecise patterns of connectivity takes place in an activity-dependent manner (for reviews see Purves & Lichtman, 1980; Shatz, 1990). In addition to activity-dependent changes in neuronal circuitry that are based on morphological alterations in connectivity, the efficacy of synaptic transmission in existing connections can also be modulated by electrical activity. Hebb (1949) has proposed that

coinciding pre- and postsynaptic activity may lead to changes in synaptic strength. Similar mechanisms are believed to underlie long-term potentiation (LTP) (for reviews see Brown *et al.*, 1990; Madison *et al.*, 1991; Tsumoto, 1992) and synaptogenesis in the developing brain (e.g. Constantine-Paton, 1990). There is also accumulating evidence for the presence of activity-dependent long-term depression (LTD) in the cerebellum (Ito, 1989), hippocampus (Stanton & Sejnowski, 1989) and neocortex (Artola *et al.*, 1990). The maturation of neuronal properties and neuronal differentiation, too, seems to be under control of electrical activity (Spitzer, 1991). For example, electrical activity might be crucial for the full maturation of inhibitory synaptic mechanisms (Hendry & Jones, 1988; Corner & Ramakers, 1992).

### 1.4. NETWORK FORMATION AND OVERSHOOT

A wide range of processes having great significance for network formation appear to be modulated by neuronal activity. As a result, a mutual influence exists between the formation of synaptic connectivity and neuronal electrical activity, i.e. a feedback loop exists between changes in network structure and changes in network activity (Von der Malsburg & Singer, 1988). This feedback loop must be expected to have major implications not only for the structure of the mature network, but also for the stages a network goes through during its development.

A general feature of nervous system development, *in vivo* as well as *in vitro*, is that virtually all structural elements show an initial overproduction, followed by an elimination during further development. These so-called overshoot phenomena occur, for example, with respect to neuron numbers (e.g. Finlay & Slatery, 1983; Heumann & Leuba, 1983; Cowan *et al.*, 1984; for reviews see Oppenheim, 1991; Ferrer *et al.*, 1992), connections (e.g. Kato *et al.*, 1985; Price & Blakemore, 1985; Stanfield & O'Leary, 1985), total dendritic length (Uylings *et al.*, 1990), number of dendrites (Miller, 1988), number of axons (Schreyer & Jones, 1988; Gorgels *et al.*, 1989), number of synapses (e.g. Purves & Lichtman, 1980; Huttenlocher *et al.*, 1982; Warren & Bedi, 1984; O'Kusky, 1985; *in vitro*: Van Huizen *et al.*, 1985, 1987), receptors (e.g. Insel *et al.*, 1990; McDonald *et al.*, 1990; for review see McDonald & Johnston, 1990), and expression of neurotransmitters (e.g. Wahle & Meyer, 1987; Parnavelas *et al.*, 1988).

In most neural network models, the activity patterns of the system are studied in response to external input, given a particular collection of cells and network structure. In developing networks, however, the number of cells and network structure are variable

and under control of the network activity itself (via the above mentioned activity-dependent processes). Furthermore, in the initial stages of development activity patterns that are not evoked by external input play a large role (see also Corner, 1990). Insight into the implications of activity-dependent processes and endogenous activity will therefore be indispensable for understanding the ontogenetic stages of the nervous system. In this article, we will address the implications of one of these processes, namely activity-dependent neurite outgrowth. It will be shown that this process can account for the occurrence of a transient overproduction of connections or synapses.

No simulation models for explaining overshoot are known to us. In Feinberg *et al.* (1990) only a descriptive statistical model is presented for some overshoot phenomena *in vivo*.

## 2. The Model

The model is not meant to mimic a particular nervous system, but is rather used as a tool to gain insight into the role of activity-dependent outgrowth for network formation in general. We use a distributed model in which initially disconnected neurons organize themselves into a synaptically connected network by neurite outgrowth and synaptogenesis, under influence of endogenous activity (there is no external input). Growing neurons are modelled as expanding neuritic fields, and the outgrowth of each neuron depends upon its own level of electrical activity. Neurons become connected when their neuritic fields overlap. All connections are taken to be excitatory, considering that the predominant form of synaptic activity during early stages of development appears to be excitatory (Jackson *et al.*, 1982; O'Brien & Fischbach, 1986; Habets *et al.*, 1987; Cherubini *et al.*, 1991; Corner & Ramakers, 1992). The model is inspired in part by tissue cultures of dissociated cerebral cortex cells (Van Huizen *et al.*, 1985, 1987; Van Huizen, 1986; Ramakers *et al.*, 1991). As in the model, cells in such cultures become organized into a network without the influence of external or sensory input.

### 2.1. NEURON MODEL

The shunting model (Grossberg, 1988; Carpenter, 1989), which mirrors the underlying physiology of simple nerve cell dynamics (Hodgkin & Huxley, 1952), is used to describe neuronal activity. In this model, excitatory inputs drive the membrane potential towards a finite maximum (or saturation potential, e.g. the  $\text{Na}^+$  equilibrium potential), while

inhibitory inputs (if any) drive the membrane potential towards a finite minimum (e.g. the  $\text{K}^+$  equilibrium potential).

For a purely excitatory network, the shunting model becomes

$$\frac{dX_i}{dt} = -X_i/\tau + (1 - X_i) \sum_j^N W_{ij} F(X_j) \quad (1)$$

where  $X_i$  is the (mean) membrane potential of neuron  $i$ ,  $F(X_j)$  is the mean firing rate of neuron  $j$ ,  $W_{ij}$  is the coupling strength between neuron  $i$  and  $j$  ( $W_{ij} \geq 0$ ;  $W_{ij}$  is defined in section 2.2), and  $N$  is the total number of neurons. Thus, eqn (1) takes the output of a neuron to be a mean firing rate. The effect of neuron  $j$  on  $X_i$  is mediated by trains of action potentials, and hence is proportional to the product of  $F(X_j)$  and  $W_{ij}$ . The sum of all excitatory inputs,  $\sum_j^N W_{ij} F(X_j)$ , drives  $X_i$  towards the excitatory saturation potential, which is set equal to 1. In the absence of inputs,  $X_i$  decays, with a rate determined by  $1/\tau$ , to the resting potential, which is set equal to 0. Thus,  $X_i$  is scaled between 0 and 1. The initial values of  $X_i$  are set below the saturation potential (i.e.  $X_i < 1$ ). The firing rate function  $F$  is a sigmoidal function of the membrane potential

$$F(X) = \frac{1}{1 + e^{(\theta - X)/\alpha}} \quad (2)$$

where  $F(X)$  is the firing rate [with its maximum set to 1, see Fig. 1(b)],  $\alpha$  determines the steepness of the function and  $\theta$  represents the firing threshold. The low firing rate when  $X$  is sub-threshold may be considered as representing spontaneous activity, arising from threshold fluctuations (Verveen, 1960), membrane potential fluctuations, synaptic noise (Korn & Faber, 1987), and/or random external input. Equation (1) and similar ones have been widely used in the field of neural network modelling (Grossberg, 1988; and references therein).

### 2.2. OUTGROWTH AND CONNECTIVITY

Neurons are randomly placed on a two-dimensional surface. In order to model outgrowth and formation of connections, each neuron is given a circular "neuritic field", the radius of which is variable. When two such neuritic fields overlap, both neurons, say  $i$  and  $j$ , become connected to each other with a strength proportional to the area of overlap

$$W_{ij} = A_{ij}c \quad (3)$$

where  $A_{ij} = A_{ji}$  is the amount of overlap ( $A_{ii} = 0$ ) and  $c$  is a constant of proportionality;  $A_{ij}$  may be regarded as representing the total number of synapses formed between neuron  $i$  and  $j$ , while  $c$  could represent the




FIG. 1. The size of the hysteresis loop [eqn (9)] depends on  $\alpha$  of the firing rate function  $F(X)$  [eqn (2)]. The smaller  $\alpha$ , the steeper  $F(X)$  (b) and the larger the hysteresis loop (a). Shown are  $\alpha = 0.12, 0.10$  and  $0.08$ .

synaptic strength. In this abstraction, no distinction has been made between axons and dendrites. If one were to do so, this would lead to a connectivity matrix  $\mathbf{W}$  that needs no longer be symmetric. In most of the simulations we used eqn (3). Just in order to test whether asymmetry would affect the results, we also used

$$W_{ij} = A_{ij}C_{ij} \quad (4)$$

where  $A_{ij}$  is as before and  $C_{ij}$  is uniformly distributed with mean  $c$ , so  $\mathbf{W}$  becomes asymmetric.

In the model, the outgrowth of each individual neuron depends upon its own level of electrical activity. Since the effect of activity on outgrowth is likely to be mediated by  $[\text{Ca}^{2+}]_{\text{in}}$  (Cohan *et al.*, 1987; Fields *et al.*, 1990b; Kater & Mills, 1991), and one of the consequences of the firing of action potentials is the influx of calcium ions (e.g. Hockberger *et al.*, 1989), leading to elevated  $[\text{Ca}^{2+}]_{\text{in}}$ , we take the outgrowth to be dependent upon the firing rate

$$\frac{dR_i}{dt} = \rho G(F(X_i)) \quad (5)$$

where  $R_i$  is the radius of the circular neuritic field of neuron  $i$ , and  $\rho$  determines the rate of outgrowth. The outgrowth function  $G$  is defined as

$$G(F(X_i)) = 1 - \frac{2}{1 + e^{(\epsilon - F(X_i))/\beta}} \quad (6)$$

where  $\epsilon$  is the value of  $F(X_i)$  for which  $G = 0$  and  $\beta$  determines the steepness of the function. The function  $G$  remains in the bounded range  $\langle -1, 1 \rangle$ . Depending on the firing rate, a neuritic field will grow out [ $G > 0$  when  $F(X_i) < \epsilon$ ], retract [ $G < 0$  when  $F(X_i) > \epsilon$ ] or remain constant [ $G = 0$  when  $F(X_i) = \epsilon$ ] (see Fig. 2). Equation (6) is simply a phenomenological description of the theory of Kater *et al.* (Kater *et al.*,

1988; Kater & Guthrie, 1990; Kater *et al.*, 1990) that the electrical activity of a neuron affects (via calcium influx) its outgrowth. High activity or  $[\text{Ca}^{2+}]_{\text{in}}$  produces retraction, as reported in (Cohan & Kater, 1986; Fields *et al.*, 1990a; Schilling *et al.*, 1991), and low activity or  $[\text{Ca}^{2+}]_{\text{in}}$  allows outgrowth. This description is also in agreement with observations that suppression of activity favours neurite outgrowth (Van Huizen & Romijn, 1987; Schilling *et al.*, 1991). An admittedly unrealistic property of eqn (6) is that if  $F(X_i) < \epsilon$ , a neuron could grow out indefinitely. We preferred, however, not to put explicit bounds on the neuritic field size, because it appeared that the network itself is capable of regulating the size of its neurons.

Essential in the formulation of the outgrowth model is that connectivity can change during development. Using growing neuritic fields and taking the connection strength proportional to the area of overlap is just one way of achieving this. In addition, we have also studied several other ways of changing connectivity (see Section 4).

To summarize, each neuron is described by differential equations for both the membrane potential  $X$  and the radius of the neuritic field  $R$ . In total, the




FIG. 2. Outgrowth function  $G$  [eqn (6)].

model thus consists of  $2N$  differential equations, where  $N$  is the total number of neurons. The connectivity matrix  $\mathbf{W}(N \times N)$  is variable and is determined by calculating the degree of overlap of the neuritic fields. The model is studied both analytically and by means of numerical solution, employing the variable time step Runge-Kutta integrator provided by Press *et al.* (1988).

### 2.3. PARAMETERS

The membrane time constant  $\tau$  was set at 8 msec, which is well within the range of values reported for cortical cells (Connors *et al.*, 1982) and hippocampal cells (Lacaille *et al.*, 1987). However, the results will appear not to depend upon the actual choice for  $\tau$ . The outgrowth of neurons is on a time scale of days or weeks (Van Huizen *et al.*, 1985, 1987; Van Huizen, 1986; Ramakers *et al.*, 1991; Schilling *et al.*, 1991), so that connectivity can be regarded as quasi-stationary on the time scale of membrane potential dynamics (i.e.  $\rho$  much smaller than  $1/\tau$ ). To avoid unnecessarily slowing down the simulations,  $\rho$  was chosen as large as possible so as to maintain the quasi-stationary approximation. In most simulations, we used  $\rho = 2.5 \times 10^{-6}$ . As nominal values for the other parameter, we choose  $\theta = 0.5$ ,  $\alpha = 0.10$ ,  $\beta = 0.10$  and  $\epsilon = 0.60$ . The effect of other values was studied in order to test the robustness of the results (see Section 4).

## 3. Results

### 3.1. GLOBAL MODEL BEHAVIOUR

The behaviour of the model can in part be predicted directly from eqn (1). The excitatory network defined by eqn (1) ( $W_{ij} \geq 0$ ;  $W_{ij}$  need not equal  $W_{ji}$ ) has, for biologically realistic initial conditions ( $X_i < 1$ ), convergent dynamics (Hirsch, 1989): every trajectory converges to some equilibrium point. The equilibrium states of the system for a given  $\mathbf{W}$  are given by the solutions of

$$0 = -X_i/\tau + (1 - X_i) \sum_j^N W_{ij} F(X_j) \quad \forall i. \quad (7)$$

If the variations in  $X_i$  are small (i.e. relative to  $\bar{X}$ ), we find for the average membrane potential of the network (see Appendix)

$$0 \simeq -\bar{X}/\tau + (1 - \bar{X}) \bar{W} F(\bar{X}) \quad (8)$$

where  $\bar{W} = (1/N) \sum_{ij}^N W_{ij}$ . By means of numerical simulation, it was confirmed that eqn (8) captures the essentials for qualitatively describing the global




FIG. 3. Hysteresis. Steady state ( $d\bar{X}/dt = 0$ ) dependence on  $\bar{W}$  ( $\bar{W} = (1/N) \sum_{ij}^N W_{ij}$ ), according to eqn (9). See further Section 3.1 of text.

behaviour of the model in terms of  $\bar{W}$  and  $\bar{X}$  (see Section 3.2). Based on this approximation,  $\bar{W}$  can be written as a function of  $\bar{X}$

$$\bar{W} = \frac{\bar{X}/\tau}{(1 - \bar{X})F(\bar{X})} \quad 0 \leq \bar{X} < 1 \quad (9)$$

which gives the steady state ( $d\bar{X}/dt = 0$ ) dependence on  $\bar{W}$  (Fig. 3). The steady states lying on the branch  $ABC$  (quiescent states:  $\bar{X}$  is low) and on  $DEF$  (activated states:  $\bar{X}$  is high) are stable, whereas those on  $CD$  are unstable. Thus for  $w_1 < \bar{W} < w_2$  ( $w_1$  and  $w_2$  are the critical points), there exist two stable steady states (see also Murray, 1989). For slowly increasing  $\bar{W}$  (starting at  $A$ ) the path followed by  $\bar{X}$  is different from that for slowly decreasing  $\bar{W}$  (starting at  $F$ ):  $ABCE$  and  $FEDBA$ , respectively. The presence of this hysteresis loop underlies the emergence of overshoot.

The size of the neuritic fields, and therefore  $\bar{W}$ , is governed by the system itself, being under control of neuronal activity. A neuritic field remains constant for  $G = 0$ , i.e. if  $X_i = F^{-1}(\epsilon)$  [eqn (6)], where  $F^{-1}$  is the inverse function of  $F$ . If this holds for all cells,  $\bar{W}$  remains constant. Then, since all cells have identical  $\epsilon$ , also  $\bar{X} = F^{-1}(\epsilon)$ . Thus, the equilibrium point of the system is the intersection point of the line  $\bar{X} = F^{-1}(\epsilon)$  with the curve of Fig. 3. The rate of outgrowth is so low relative to the dynamics of the membrane potential that, for changing  $\bar{W}$ ,  $\bar{X}$  follows the curve. Starting with initially disconnected cells ( $\bar{W} = 0$ ), the model can display four qualitatively different kinds of global behaviour (see Fig. 4), depending on the position of the equilibrium point (on branch  $DE$ ,  $EF$ ,  $ABC$ , or  $CD$ ). Those lying on the branch  $CD$  are unstable, the rest are stable.




FIG. 4. Behaviour of the model (with  $c = 0.1$ ,  $N = 64$ ) for different values of  $\epsilon$ . Connectivity  $C$ , against time, and against average membrane potential  $\bar{X}$  ( $C = \frac{1}{2} \sum_{ij} A_{ij}$  = total area of overlap =  $N\bar{W}/2c$ ). (a)  $\epsilon = 0.60$ : overshoot in connectivity; (b)  $\epsilon = 0.97$ : no overshoot; (c)  $\epsilon = 0.30$ : oscillations.

#### *DE: overshoot*

Because the activity in the network is initially low,  $dR_i/dt$  is positive and  $\bar{W}$  increases, whereby  $\bar{X}$  follows the branch  $ABC$  until it reaches  $w_2$ , at which point  $\bar{X}$  jumps to the upper branch, thus exhibiting a phase transition from quiescent to activated state. The activity in the network is then however so high [i.e. above  $F^{-1}(\epsilon)$ ], that the neuritic fields begin to retract ( $dR_i/dt < 0$ ) and  $\bar{W}$  to decrease, whereby  $\bar{X}$  moves along the upper branch from  $E$  to the intersection point. Thus, in order to arrive at an equilibrium point on the branch  $DE$ , a developing network has to go through a phase in which  $\bar{W}$  is higher than in the final situation, thus exhibiting a transient overshoot in  $\bar{W}$ . Recall that  $\bar{W}$  represents the number of synapses or connections in the network.

In Section 4, it will be shown that the existence of a hysteresis loop hinges upon the firing rate function  $F$  having a firing threshold and low but non-zero values for sub-threshold membrane potentials. The size of the hysteresis loop depends on  $\alpha$  (Fig. 1). A

smaller  $\alpha$  results in a larger hysteresis loop and a larger overshoot in  $\bar{W}$ . Since for small  $\bar{X}$  the increase of  $\bar{X}$ , starting with  $\bar{W} = 0$  and  $\bar{X} = 0$ , is proportional to  $\bar{W}F(\bar{X})$  [see eqn (1)], and the increase in  $F(\bar{X})$  is relatively slow due to  $F$  being concave for  $X < \theta$ ,  $\bar{X}$  increases mainly as the result of increasing  $\bar{W}$ . Thus, the lower the value of  $F(X)$  for sub-threshold values of  $X$  (i.e.  $\alpha$  small), the higher  $\bar{W}$  should become to activate the network, resulting in a larger hysteresis loop [if  $F(X) = 0$  for low  $X$ , one would need infinitely high  $\bar{W}$  to activate the network]. Once  $F(\bar{X})$  is high,  $\bar{W}$  may be lower to keep  $\bar{W}F(\bar{X})$  high. In other words, a higher  $\bar{W}$  is needed to trigger activity in a quiescent network than to sustain it once the network has been activated.

#### *ABC and EF: no overshoot*

If the neuritic fields start retracting already at a very low level of activity, or only when the level of activity is very high, no overshoot occurs. In the first case, the equilibrium point lies on  $ABC$ , and the

neurons stop growing even before  $\bar{W}$  can reach the critical point  $w_2$ . In the second case, the equilibrium point lies on  $EF$ , and  $\bar{W}$  will remain increasing after  $\bar{X}$  has jumped to the upper branch, which is followed until the intersection point is reached.

*CD: oscillations*

An intersection point on this branch is unstable and results in regular oscillations that follow the path  $ABCEDBCEDBC\dots$ . The period of these oscillations is determined by the value of  $\rho$ .

3.2. SIMULATION RESULTS

*Approximation*

The validity of approximating the global behaviour of the network by eqn (9) was confirmed by numerical simulation (Fig. 4). Generally, it was found that this approximation is good whenever the synaptic strength  $c$  is such that, at the time that (part of) the network approaches the transition to the activated state, the connectivity matrix  $\mathbf{W}$  is a connected graph, i.e. there is a path from each cell to any other cell (see Peretto & Niez, 1986). This can be achieved when




FIG. 5. Neuritic field sizes. (a) Early stage of development. The same network as in Fig. 6. The numbers correspond to those in Fig. 6. (b) The same network as in (a) at equilibrium. Neuritic field size depends on local cell density. (c) Network ( $N = 25$ ,  $c = 0.2$ ) at equilibrium with cells at grid positions. (d) Network ( $N = 25$ ,  $c = 0.4$ ) at equilibrium with cells at noisy grid positions. (e) Network ( $N = 25$ ) with high synaptic strength ( $c = 2.5$ ), at equilibrium. Cells develop into sub-networks. (f) Same network as in (e) but with  $c = 0.3$ .

$c$  is low enough that for the network to become activated each cell has to be connected to a number of neighbouring cells. If  $c$  is too high, on the other hand, the network breaks up into sub-networks [see Fig. 5(e)], in which the transition to the activated state may take place at different times, depending on the local cell density. The phase transition of the whole network is then fragmented and less clearcut: in some parts of the network the cells may already be retracting, while in others they are still growing out. [In Fig. 4(a) the relative overshoot is larger than in Fig. 6 because  $c$  is smaller.] The approximation is then no longer valid. Initially, eqn (8) is always a good approximation, because development starts with disconnected cells. At equilibrium, the network is exactly described by eqn (8), since then  $X_i = F^{-1}(\epsilon)$ ,  $\forall i$ .

*Local behaviour*

Except for their position, all the cells are exactly identical. Local variations in cell density, however, suffice to generate a great variability among individual cells, with respect both to their neuritic field size at equilibrium [Fig. 5(b), (d)–(f)] and to their developmental course of field size and firing behaviour (Fig. 6).

All cells will attain a fixed equilibrium size for which the input from overlapping cells is such that  $F(X_i) = \epsilon$ ,  $\forall i$ . Cells surrounded by a high number of neighbouring cells tend to become small since a small

neuritic field will already give sufficient overlap with other cells. In contrast, relatively isolated cells must grow large neuritic fields in order to contact a sufficient number of cells. One might say that the neuritic fields adapt to the available space so as to cover it optimally. Because all cells have the same firing rate function and  $\epsilon$ , the area of overlap with other cells,  $\sum_j^N W_{ij}$ , will be the same for all cells. This area is given by [using eqn (6) and eqn (7)]

$$\sum_j^N W_{ij} = \frac{F^{-1}(\epsilon)/\tau}{\epsilon(1 - F^{-1}(\epsilon))} \tag{10}$$

In dense parts of the network, the transition to the activated state takes place earlier than in less dense parts. Interaction between areas that are, and those that are not yet activated causes a range of developmental patterns in firing behaviour. A cell bordering on a cluster of cells with which it has contact at the moment the cluster makes a phase transition, will show a pattern like Fig. 6.3, in which the subsequent relatively rapid decrease in activity within this cluster, followed by further outgrowth of the border cell, gives rise to a damped oscillation in firing rate. An isolated cell making contact with a cluster only after this cluster has gone through its phase transition (and to a large extent completed its subsequent decrease in field sizes and activity levels), will show a pattern like Fig. 6.6. Since the activity within the cluster remains essentially constant, this cell will continue to grow out




FIG. 6. Local behaviour. Same network as in Fig. 4, but with  $c = 0.4$ , and  $\epsilon = 0.6$ .  $C = \frac{1}{2} \sum_j^N A_{ij}$  = total area of overlap =  $NW/2c$ . In the first row the average behaviour of the network is shown; 1, . . . , 6 are individual cells. Note that the relative overshoot in  $R$  is less than in  $C$  because (i)  $C$  is an area and  $R$ , a radius; and (ii)  $R_i$  contributes to  $C$  only if the neuritic fields actually overlap.



until its overlap is such that  $F(X_i) = \epsilon$ . In contrast, a cell at the heart of such a cluster will display a pattern like Fig. 6.1.

*Network size*

Synaptic strength can determine the size of the resulting network(s) [Fig. 5(e) and (f)]. When  $c$  is low, cells develop into one, connected network, whereas a high  $c$  can result in a development into separate networks, because contact with fewer cells is sufficient for  $F(X_i) = \epsilon$ .

*Timing of overshoot*

In networks with a high density of cells, overshoot takes place earlier than in low-density networks (where neurons have to grow for a longer period of time in order to make sufficient contacts). Also the higher the synaptic strength  $c$ , the earlier the over-

shoot will occur. If activity is totally blocked, on the other hand, cells keep growing out, and no reduction in connectivity can take place at all.

**4. Robustness**

The robustness of the results was tested under different parameter values and alternative formulations of the model.

**4.1. FIRING RATE FUNCTION**

The smaller  $\alpha$  in eqn (2) is, the steeper  $F$  and the larger the hysteresis loop will be, resulting in a more pronounced overshoot (see Fig. 1 and section 3.1). Lowering  $\theta$ , which translates  $F$  to the left, also increases the firing rate at low  $X$ . Thus, without simultaneously lowering  $\alpha$ , this would result in a smaller hysteresis loop. In Fig. 7(h), a firing rate function is used with  $\theta = 0.2$  and  $\alpha = 0.05$ : In fact,  $F$




FIG. 7. Different firing rate functions (firing rate  $F$  versus membrane potential  $X$ ) together with the curve defined by eqn (9) ( $\bar{X}$  versus  $\bar{W}$ ). Except for linear and convex functions, all firing rate functions give rise to a hysteresis loop. (a) Piece-wise linear. (b) Discrete threshold ( $F = 0.01$  for  $X < 0.5$ ). (c)  $F = (1.0 - s)X + s$ ;  $s = 0.01$ . (d)  $F = X^2 + s$ ;  $s = 0.0025$ . (e)  $F = X/(X + K) + s$ ;  $K = 0.25$ ,  $s = 0.01$ . (f) Sigmoidal function based on Michaelis-Menten function (De Boer & Perelson, 1991).  $F = (1 - s)X^2/(\theta^2 + X^2) + s$ ;  $\theta = 0.2$ ,  $s = 0.001$ . (g) Composite function. For  $X \leq 0.3$ ,  $F = 0.5/(1 + e^{(\theta - X)/\alpha})$  with  $\alpha = 0.055$  and  $\theta = 0.2$ . For  $X > 0.3$ ,  $F$  is linear. (h) Sigmoidal function according to eqn (2), with  $\alpha = 0.05$  and  $\theta = 0.2$ .

in eqn (1) may be replaced by any other function so long as the first part has a positive first and second derivative (or a discrete threshold) and  $F(X) > 0$  for small  $X$  including  $X = 0$ . All these functions will give rise to hysteresis (Fig. 7). Note that for overshoot to occur it is not essential that the firing rate during the phase transition jumps to a high value relative to its maximum, as it does with eqn (2). Provided the value of  $\epsilon$  is changed accordingly, a function like that of Fig. 7(g) can also be used.

#### 4.2. NEURON MODEL

As can be seen from eqn (9),  $\tau$  is only a scale factor and has no influence on the shape of the hysteresis curve.

The general result is also obtained in some alternative formulations of the neuron model. For example, in the additive model (Grossberg, 1988)

$$\frac{dX_i}{dt} = -X_i/\tau + \sum_j^N W_{ij}F(X_j) \quad (11)$$

which lacks the factor  $(1 - X_i)$  in eqn (1), so  $X_i$  is no longer bounded. In analogy with eqn (9) we obtain,

$$\bar{W} = \frac{\bar{X}/\tau}{F(\bar{X})} \quad (12)$$

which is plotted in Fig. 8(a) [with  $F$  according to eqn (2)]. Because  $F$  is a sigmoidal, saturating function,  $1 - X_i$  may be omitted without losing hysteresis. If, on the other hand,  $F$  is without bound (e.g. a power function) the saturating factor  $1 - X_i$  is necessary for hysteresis. A modified form of eqn (1) was introduced by Wilson & Cowan (1972)

$$\frac{dX_i}{dt} = -X_i/\tau + (1 - X_i)F\left(\sum_j^N W_{ij}X_j\right) \quad (13)$$

which replaces the sum of non-linear signals in eqn (1) by a non-linear function of the sum. In analogy with eqn (9) we obtain,

$$F(\bar{W}\bar{X}) = \frac{\bar{X}/\tau}{1 - \bar{X}} \quad (14)$$

and, using eqn (2)

$$\bar{W} = (\theta - \alpha \ln(\tau/\bar{X} - \tau - 1))/\bar{X} \quad 0 < \bar{X} < \frac{\tau}{1 + \tau} \quad (15)$$

which is plotted in Fig. 8(b).

#### 4.3. NETWORK SIZE

This is not a crucial parameter. The same phenomena are retained, for example, in a two-neuron model

$$\begin{aligned} \frac{dX_1}{dt} &= -X_1/\tau + (1 - X_1)WF(X_2) \\ \frac{dX_2}{dt} &= -X_2/\tau + (1 - X_2)WF(X_1). \end{aligned} \quad (16)$$

The conditions for hysteresis and overshoot can now be formulated in terms of intersecting isoclines  $dX_1/dt = dX_2/dt = 0$  (Fig. 9). In order for hysteresis to occur there must be three equilibrium points (two stable and one unstable) for intermediate values of  $W$ . This can be achieved by sigmoidal isoclines (the form of the isoclines is determined by  $F$  and  $1 - X_i$ ).

Even a single neuron that is connected only to itself (Segal & Furshpan, 1990) is capable of displaying overshoot; in such a case the null-isocline is exactly eqn (8).

#### 4.4. OUTGROWTH FUNCTION

Parameter  $\beta$  in eqn (6) determines the steepness of  $G$ . The form of the outgrowth function, however, affects only the specific time course of neurite outgrowth. In fact, any other function  $G$  for which  $G > 0$  at low values of  $F(X_i)$  and  $G < 0$  at high values of




FIG. 8. Steady-state dependence ( $d\bar{X}/dt = 0$ ) on  $\bar{W}$  for (a) additive model eqn (11) and (b) model according to eqn (13).




FIG. 9. Isoclines of the two-neuron model for (a) a low, (b) an intermediate and (c) a high value of  $W$ . Arrows in (b) indicate the direction of the trajectories. For intermediate values of  $W$ , there exist three steady states, one of which (the middle one) is unstable.

$F(X_i)$  may be used. The same results are also obtained if outgrowth is made directly dependent on  $X_i$ , instead of on  $F(X_i)$ . The firing rate at which a neuritic field stabilizes,  $\epsilon$ , is an especially important parameter of the model in that it determines whether or not overshoot will occur, and, if so, what its relative size will be (see Fig. 3).

#### 4.5. CONNECTIVITY

Whether the connectivity matrix  $W$  is symmetric or asymmetric does not play a role in eqn (9). This implies that modelling axons and dendrites separately would not affect the main findings. For example, using eqn (4), with  $C_{ij}$  uniformly distributed between 0.0 and 0.8, yielded similar results as in the experiment of Fig. 6. The presence of hysteresis and overshoot also does not depend on the particular way in which connections between cells are defined. In Fig. 10, the results are shown of an experiment in which  $A_{ij}$  of eqn (3) is simplified to  $A_{ij} = 1$  if the neuritic fields of neuron  $i$  and  $j$  overlap, and  $A_{ij} = 0$  if they do not. Even ignoring spatial structure by randomly filling in  $W$  gives the same general outcome.




FIG. 10. The average membrane potential against the average number of connections. Number of connections  $= (1/N) \sum_{i,j}^N A_{ij}$ , in an experiment where  $A_{ij}$  of eqn (3) is simplified to  $A_{ij} = 1$  if neuron  $i$  and  $j$  overlap, and  $A_{ij} = 0$  if they do not. With  $N = 25$ ,  $c = 0.05$ ,  $\epsilon = 0.65$  and  $\rho = 5.0 \times 10^{-5}$ .

#### 4.6. VARIATION

The results are robust with respect to differences in intrinsic properties, e.g. by randomly drawing cell parameters from uniform distributions. Thus, with  $\alpha$  uniformly distributed over  $[0.08, 0.12]$ ,  $\tau \sim U[7, 10]$ ,  $\theta \sim U[0.4, 0.6]$ ,  $\epsilon \sim U[0.6, 0.8]$ ,  $\beta \sim U[0.08, 0.12]$ , and  $c = 0.1$  and  $N = 64$ , similar results were obtained as in Fig. 4. Thus, overshoot can also occur in networks composed of different cell types.

Although the connectivity structure of the final, stable network is constrained by the fact that each cell must have inputs from overlapping cells such that  $F(X_i) = \epsilon$ ,  $\forall i$ , no unique solution exists for a given position of the cells. The initial conditions ( $R_i$ 's) and the other cell parameters, which may be different for different cells, will determine which network of all possible ones is actually built.

A network in which  $\epsilon$  is distributed over a range of values that includes unstable ones, can still show overshoot as a whole, although the dynamics of the individual cells may be very complex. We are currently investigating such possibilities.

#### 4.7. EXTERNAL INPUT

The neuron model with external input becomes

$$\frac{dX_i}{dt} = -X_i/\tau + (1 - X_i) \left( E_i + \sum_j^N W_{ij} F(X_j) \right) - (1 + X_i) I_i \quad (17)$$

where  $E_i$  and  $I_i$  are external excitatory and external inhibitory input, respectively; the inhibitory saturation potential is set to  $-1$ . In analogy with eqn (9) we obtain

$$\bar{W} = \frac{\bar{X}/\tau - (1 - \bar{X})E + (1 + \bar{X})I}{(1 - \bar{X})F(\bar{X})} \quad -1 < \bar{X} < 1 \quad (18)$$

which is plotted in Fig. 11 for ( $E = 0.015, I = 0$ ), ( $E = I = 0$ ; normal situation), and ( $I = 0.008, E = 0$ ). Without adjusting  $\alpha$ , external excitatory input makes




FIG. 11. Effect of external input. External excitatory input makes the hysteresis loop smaller, while inhibitory input makes it larger. Eqn (18) for  $E=0.015$ ,  $I=0$ ;  $E=I=0$  (normal situation), and  $I=0.008$ ,  $E=0$ .

the hysteresis loop smaller, while inhibitory input makes it larger. Thus, with excitatory input,  $\alpha$  can be much lower—or even  $F(X)$  such that  $F(X)=0$  for  $X$  below a certain value—and still not result in an excessively large hysteresis loop.

### 5. Comparison with Empirical Data

The results show some striking similarities, with respect to development of electrical activity, cell morphology, and connectivity, with what has been observed in developing *in vitro* cultures of dissociated cells.

The sequence of events in the model, with an initial phase of neurite outgrowth while electrical activity is low, an abrupt transition to high activity when connectivity reaches a critical value, and a phase with neurite retraction thereafter, has also been observed in cerebellar cultures (Schilling *et al.*, 1991). Purkinje cell dendrites elongate steadily during the first week after plating, when electrical activity is still negligible. As a result of cells becoming integrated into a functional neuronal network, electrical activity increases dramatically between 7–10 days *in vitro*. During this transition period dendrites cease their linear growth, retract, and then begin to branch profusely (in our model implementation, using neuritic fields, no separate branching events are distinguished). Coinciding with this period, there is an increase in  $[Ca^{2+}]_{in}$ . The notion that electrical activity, presumably by enhancing  $[Ca^{2+}]_{in}$ , might regulate dendritic growth patterns was further supported by the observation that blockade of electrical activity resulted in the continued elongation of dendrites as well as a decrease in  $[Ca^{2+}]_{in}$ . The developmental sequence described above applied to all Purkinje

cells, but the actual timing varied for individual cells. Variations in timing are also observed in the model, and occur simply as a result of differences in local density. The model cells are capable at any time of changing the length of their neurites in response to changes in activity. Morphological changes in Purkinje cells, too, were found to be reversible, with even mature-looking cells altering their dendritic growth in response to changes in activity. Schilling *et al.* (1991) propose that in the absence of synaptic input dendrites will continue to elongate until they encounter active presynaptic elements. This view is consistent with our model, in which a disconnected cell will grow out until it has established sufficient contacts with active cells (see Section 3.2). It is also consistent with the synaptotrophic hypothesis of dendritic growth (Vaughn, 1989), which posits that growth and branching are initiated and maintained by synapses formed on growth cones.

As in the simulation model, cultures of dissociated cerebral cortex cells show a transient overproduction (overshoot) of synapses during development (Van Huizen *et al.*, 1985, 1987; Van Huizen, 1986), with a phase of neurite outgrowth and synapse formation during the first 3 weeks *in vitro* being followed by a substantial elimination of synapses during the week thereafter (Fig. 12). The development of electrical activity in these cultures shows a good correspondence with what has been described above for cerebellar cultures (and for the simulation model). With increasing synaptic density, single neuron firing and network activity—which takes the form of repetitive slow field potentials (Corner & Crain, 1972; Van Ooyen *et al.*, 1992)—abruptly appear within a window of a few days (Habets *et al.*, 1987). In such cultures, electrical activity appears to control both neurite outgrowth and synapse elimination: chronic blockade of electrical activity enhanced neurite outgrowth (Van Huizen & Romijn, 1987) and prevented synapse elimination during the fourth week (Van Huizen *et al.*, 1985). In contrast, chronic intensification of activity accelerated the formation of synapses and hastened the process of synapse elimination by almost a week (Van Huizen *et al.*, 1987) (Fig. 12). The model responds in a similar way to suppression or intensification of electrical activity (see Section 3.2, *Timing of overshoot*).

Different cell types in the rat cerebral cortex show different growth curves with respect to radial distance and total dendritic length (Uylings *et al.*, 1990), with cells displaying clear, minor or no overshoot at all. Some of these curves show a clear resemblance to the ones observed in the model.




FIG. 12. Cultures of dissociated cerebral cortex cells show a transient overproduction of synapses (control). Chronic blockade of activity (by TTX) largely prevents synapse elimination, whereas intensification (PTX) accelerates the process. After Van Huizen (1986).

## 6. Conclusions and Discussion

Many of the processes that play a role in network development are dependent upon the electrical activity of the neurons and network itself, so that a tight coupling exists between network formation and network activity. In this study, we have examined the consequences for network ontogeny of one of these processes, namely activity-dependent neurite outgrowth. We have demonstrated that the very presence of this mechanism in combination with another elementary neuronal property—a non-linear neuron response function, i.e. a threshold for action potential generation—is sufficient to generate overshoot phenomena with respect to connectivity. The results are robust under: different firing rate functions (provided they have a type of firing threshold and low but non-zero values for sub-threshold membrane potentials, i.e. spontaneous activity); variance among neurons in all parameters; different neurite outgrowth functions (provided a high level of electrical activity results in retraction and a low level in outgrowth); symmetric versus asymmetric connectivity matrix; the way in which connections are defined; network size and different neuron models. The question of whether other processes such as trophic action and inhibition might interfere with the generation of overshoot was not addressed in this study. At present we are studying the effects of inhibitory neurons on network development, the preliminary results of which show that the general findings of this article remain completely true for networks with inhibitory circuits.

The mechanism underlying the generation of overshoot in this simple model may provide part of

the explanation for overshoot phenomena in the developing nervous system, at least for those which have been observed in tissue cultures of dissociated cells. In such cultures, the sequence of events corresponds closely to that observed in the simulation model.

It should be emphasized that other, more realistic, descriptions of neurite outgrowth and synaptogenesis (e.g. with individual neurites instead of a circular neuritic field, and possibly some form of conservation of, or limit upon, the total numbers of synapses per cell rather than simply assuming that the larger the overlap of neuritic fields, the more synapses will be made) are merely different mappings onto the connectivity matrix  $W$ . Although this could affect the particular network structure, overshoot would still take place.

The neuritic field size of the model cells adapts to the local cell density, resulting in small fields in dense areas and larger ones in sparse areas. In this respect, it is interesting that the dendritic fields of ganglion cells, the cell bodies of which are arrayed in a regular mosaic, have been reported to achieve a uniform coverage of the retina so that every point of the visual space is "seen" by at least one cell (Wässle *et al.*, 1981). From these observations it was inferred that some kind of growth mechanism must exist whereby local interactions among cells regulate dendritic field size. Activity-dependent outgrowth may be a candidate for such a mechanism,

Hysteresis may provide a mechanism by which overshoot phenomena in general can be understood. To illustrate this, consider the following variant of the model. Suppose that cell death is more likely at (very) high levels of electrical activity, and that neurons

(whose neuritic fields are now taken to be constant) are produced at a given source, transported and incorporated into a network during development. As before, neurons become connected when their neuritic fields overlap. At a given cell density, the network will become activated and start losing cells if the resulting activity is too high. Because of hysteresis, a higher cell density is necessary to activate the network than to sustain it once activity has been initiated, thus giving rise to a reduction in cell number. Such a transient overproduction of cells has been observed in the development of, for example, the suprachiasmatic nucleus (Swaab *et al.*, 1990). Another variant of the model would be one in which the neuritic fields are more or less constant and (partly) overlapping, while it is now the formation of new synapses or the strengthening of older ones which is activity-dependent. This would lead to exactly the same results, with overshoot in number of synapses or synaptic strength, respectively.

For some values of  $\epsilon$ , the firing rate at which a given neuritic field remains constant, the model can generate sustained oscillations in overall activity (and connectivity). The period of these oscillations is determined by  $\rho$ , the rate of outgrowth of neuritic fields or, in the alternative formulations of the model, the rate at which synapses are formed/destroyed or existing ones are strengthened/weakened. Since these changes can occur on a time scale of hours, this mechanism might provide a possible explanation for the occurrence of slow rhythmic activity in various brain areas (for example, circadian rhythms in the suprachiasmatic nucleus).

With respect to overshoot in connectivity, there may exist a parallel with the developing immune system. In the immune system, which may function as a network (Jerne, 1974), a high idiotypic connectivity (the number of clones with which a given clone interacts) is found during early ontogeny (Holmberg *et al.*, 1986). A possible explanation might be that highly connected clones, which become over-stimulated or suppressed if clones are large, can only be maintained during early life when all clones are small and the network has not yet filled with antibodies (De Boer & Perelson, 1991). In analogy, neurons in our network can only maintain a high number of connections during early development when the network is not yet "filled" with activity. Once the network is activated, these neurons become over-stimulated and subsequently lose some of their connections.

Overshoot phenomena may also be involved in learning. Doyle *et al.* (1992) presented evidence to suggest that information storage during learning may be based on connectivity changes mediated by a

replay of early developmental events. In this concept, information acquisition may induce a transient overproduction of synaptic contacts in a given network, followed by an activity-dependent selection to yield a new circuitry.

The present study might also have relevance for understanding the ontogenetic origins of epilepsy. If, for whatever reason, activity is blocked during development, insufficient reduction in connectivity will result. This could lead in adulthood, assuming that pruning of connections is largely restricted to a "critical period", to a network prone to epileptic-like activity because of its abnormally high degree of connectivity. Interestingly, Ben-Ari & Represa (1990) have suggested that use-dependent sprouting may play a role in epilepsy.

Damaging a proportion of the model cells results in increased neurite outgrowth of the remaining cells until all the cells have the same activity level as before. This is reminiscent of what happens following early prefrontal cortex lesions in rats. In such animals, an increase in dendritic arborization is found—possibly accompanied by synapse formation—which seems to correlate with the occurrence of behavioural sparing of function (Kolb & Gibb, 1991).

If the level of spontaneous activity is too low (or the synapses too weak) for the network to become activated by its intrinsic activity, neurites would continue growing out until at some stage during development the network may become activated by external input (e.g. sensory input). This could then result in retraction of neurites, the network being hyperactive due to its high degree of connectivity. Such a scenario would also give rise to overshoot. Age-related decreases in spine density and in total length of dendrites have indeed been reported for the development of Purkinje cells *in vivo* (Pentney, 1986). It may be hypothesized that changes in levels of electrical activity play a role in these morphological alterations.

The intracellular calcium concentration ( $[Ca^{2+}]_i$ ) is likely to be one of the primary variables controlling neurite outgrowth. One of the factors that can change  $[Ca^{2+}]_i$ , and hence neurite outgrowth, is electrical activity, which, in turn, is dependent upon network connectivity and network dynamics. Therefore, the mechanisms regulating  $[Ca^{2+}]_i$  may be major determinants of entire patterns of neuronal circuitry, as was suggested by Kater *et al.* (1988). Indeed, with the help of a relatively simple model of a developing neural network we have shown that activity-dependent neurite outgrowth can have a profound effect on network structure during development in that it generates pronounced overshoot phenomena with respect to connectivity.

We thank M. A. Corner, F. H. Lopes da Silva and B. Nienhuis for critically reading the manuscript, and M. P. van Veen for assistance with the figures.

## REFERENCES

- ARTOLA, A., BRÖCHER, S. & SINGER, W. (1990). Different voltage-dependent thresholds for inducing long-term depression and long-term potentiation in slices of rat visual cortex. *Nature, Lond.* **347**, 69–72.
- BEN-ARI, Y. & REPRESA, A. (1990). Brief seizure episodes induce long-term potentiation and mossy fibre sprouting in the hippocampus. *Trends Neurosci.* **13**, 312–317.
- BRENNEMAN, D. E. & EIDEN, L. E. (1986). Vasoactive intestinal peptide and electrical activity influence neuronal survival. *Proc. natn. Acad. Sci. USA.* **83**, 1159–1162.
- BROWN, T. H., KAIRISS, E. W. & KEENAN, C. L. (1990). Hebbian synapses: biophysical mechanisms and algorithms. *A. Rev. Neurosci.* **13**, 475–511.
- CARPENTER, G. A. (1989). Neural network models for pattern recognition and associative memory. *Neural Networks* **2**, 243–257.
- CHERUBINI, E., GALARSA, J. L. & BEN-ARI, Y. (1991). GABA: an excitatory transmitter in early postnatal life. *Trends Neurosci.* **14**, 515–519.
- COHAN, C. S. & KATER, S. B. (1986). Suppression of neurite elongation and growth cone motility by electrical activity. *Science* **232**, 1638–1640.
- COHAN, C. S., CONNOR, J. A. & KATER, S. B. (1987). Electrically and chemically mediated increases in intracellular calcium in neuronal growth cones. *J. Neurosci.* **7**, 3588–3599.
- CONNORS, B. W., GUTNICK, M. J. & PRINCE, D. A. (1982). Electrophysiological properties of neocortical neurons *in vitro*. *J. Neurophysiol.* **48**, 1302–1320.
- CONSTANTINE-PATON, M. (1990). NMDA receptor as a mediator of activity-dependent synaptogenesis in the developing brain. In: *Cold Spring Harbor Symposia on Quantitative Biology*, Vol. LV. pp. 431–443. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- CORNER, M. A. (1990). Brainstem control of behavior: ontogenetic aspects. In: *Brainstem Mechanisms of Behavior* (Klemm, W. R. & Vertes, R. P., eds) pp. 239–266. New York: John Wiley & Sons.
- CORNER, M. A. & CRAIN, S. M. (1972). Patterns of spontaneous bioelectric activity during maturation in culture of fetal rodent medulla and spinal cord tissues. *J. Neurobiol.* **3**, 25–45.
- CORNER, M. A. & RAMAKERS, G. J. A. (1992). Spontaneous firing as an epigenetic factor in brain development—physiological consequences of chronic tetrodotoxin and picrotoxin exposure on cultured rat neocortex neurons. *Dev. Brain Res.* **65**, 57–64.
- COWAN, W. M., FAWCETT, J. W., O'LEARY, D. D. M. & STANFIELD, B. B. (1984). Regressive events in neurogenesis. *Science* **225**, 1258–1265.
- DE BOER, R. J. & PERELSON, A. S. (1991). Size and connectivity as emergent properties of a developing immune network. *J. theor. Biol.* **149**, 381–424.
- DOYLE, E., NOLAN, P., BELL, R. & REGAN, C. M. (1992). Neurodevelopmental events underlying information acquisition and storage. *Network* **3**, 89–94.
- FEINBERG, I., THODE, H. C., CHUGANI, H. T. & MARCH J. D. (1990). Gamma distribution model describes maturational curves for delta wave amplitude, cortical metabolic rate and synaptic density. *J. theor. Biol.* **142**, 149–161.
- FERRER, I., SORIANO, E., DEL RIO, J. A., ALCANTARA, S. & AULADELL, C. (1992). Cell death and removal in the cerebral cortex during development. *Prog. Neurobiol.* **39**, 1–43.
- FIELDS, R. D., NEALE, E. A. & NELSON, P. G. (1990a) Effects of patterned electrical activity on neurite outgrowth from mouse neurons. *J. Neurosci.* **10**, 2950–2964.
- FIELDS, R. D., GUTHRIE, P. B., NELSON, P. G. & KATER, S. B. (1990b). Calcium homeostatic capacity is regulated by patterned electrical activity in growth cones of mouse DRG neurons. *Soc. Neurosci. Abstr.* **16**, 457.
- FIELDS, R. D. & NELSON, P. G. (1992). Activity-dependent development of the vertebrate nervous system. *Int. Rev. Neurobiol.* **34**, 133–214.
- FINLAY, B. L. & SLATTERY, M. (1983). Local differences in the amount of early cell death in neocortex predict adult local specializations. *Science* **219**, 1349–1351.
- GORGELS, T. G. M. F., DE KORT, E. J. M., VAN AANHOLT, H. T. H. & NIEUWENHUIS, R. (1989). A quantitative analysis of the development of the pyramidal tract in the cervical spine cord in the rat. *Anat. Embryol.* **179**, 377–385.
- GROSSBERG, S. (1988). Nonlinear neural networks: principles, mechanisms, and architectures. *Neural Networks* **1**, 17–61.
- HABETS, A. M. M. C., VAN DONGEN, A. M. J., VAN HUIZEN, F. & CORNER, M. A. (1987). Spontaneous neuronal firing patterns in fetal rat cortical networks during development *in vitro*: a quantitative analysis. *Exp. Brain Res.* **69**, 43–52.
- HAMBURGER, V. & OPPENHEIM, R. W. (1982). Naturally occurring neuronal death in vertebrates. *Neurosci. Comment* **1**, 38–55.
- HEBB, D. O. (1949). *The Organization of Behaviour: a Neuro-psychological Theory*. New York: John Wiley & Sons.
- HENDRY, S. H. C. & JONES, E. G. (1988). Activity dependent regulation of GABA expression in the visual cortex of adult monkeys. *Neuron* **1**, 701–712.
- HEUMANN, D. & LEUBA, G. (1983). Neuronal death in the development and aging of the cerebral cortex of the mouse. *Neuropath. appl. Neurobiol.* **9**, 297–311.
- HIRSCH, M. W. (1989). Convergent activation dynamics in continuous time networks. *Neural Networks* **2**, 331–349.
- HOCKBERGER, P. E., TSEN, H.-Y. & CONNOR, J. A. (1989). Fura-2 measurements of cultured rat Purkinje neurons show dendritic localization of Ca<sup>2+</sup> influx. *J. Neurosci.* **9**, 2272–2284.
- HODGKIN, A. L. & HUXLEY, A. F. (1952). A quantitative description of membrane current and its application to conduction and excitation in nerve. *J. Physiol.* **117**, 500–544.
- HOLMBERG, D., WENNERSTROM, G., ANRADE, L. & COUTINHO, A. (1986). The high idiotypic connectivity of “natural” newborn antibodies is not found in the adult mitogen-reactive B cell repertoires. *Eur. J. Immunol.* **16**, 82–87.
- HUTTENLOCHER, P. R., DE COURTEN, C., GAREY, L. J. & VAN DER LOOS, H. (1982). Synaptogenesis in human visual cortex—evidence for synapse elimination during normal development. *Neurosci. Lett.* **33**, 247–252.
- INSEL, T. R., MILLER, L. P. & GELHARD, R. E. (1990). The ontogeny of excitatory amino acid receptors in rat forebrain I: NMDA and quisqualate receptors. *Neuroscience* **35**, 31–43.
- ITO, M. (1989). Long-term depression. *A. Rev. Neurosci.* **12**, 85–102.
- JACKSON, M. B., LECAR, H., BRENNEMAN, D. E., FITZGERALD, S. & NELSON, P. (1982). Electrical development in spinal cord cell culture. *J. Neurosci.* **2**, 1052–1061.
- JERNE, N. K. (1974). Towards a network theory of the immune system. *Ann. Immunol. (Inst. Pasteur)* **125**, 373–389.
- KATER, S. B., MATTSON, M. P., COHAN, C. & CONNOR, J. (1988). Calcium regulation of the neuronal growth cone. *Trends Neurosci.* **11**, 315–321.
- KATER, S. B. & GUTHRIE, P. B. (1990). Neuronal growth cone as an integrator of complex environmental information. *Cold Spring Harbor Symposia on Quantitative Biology*, Volume LV, pp. 359–370. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- KATER, S. B., GUTHRIE, P. B. & MILLS, L. R. (1990). Integration by the neuronal growth cone: a continuum from neuroplasticity to neuropathology. In: *Progress in Brain Research*, Vol. 86, *Molecular and Cellular Mechanisms of Neuronal Plasticity in Normal Aging and Alzheimer's Disease* (Coleman, P. D., Higgins, G. A., & Phelps, C. H., eds) pp. 117–128.
- KATER, S. B. & MILLS, L. R. (1991). Regulation of growth cone behaviour by calcium. *J. Neurosci.* **11**, 891–899.

- KATO, T., HIRANO, A., KATAGIRI, T. & SASAKI, H. (1985). Transient uncrossed corticospinal fibres in the newborn rat. *Neuropathol. appl. Neurobiol.* **11**, 171-178.
- KOIKE, T. & TANAKA, S. (1991). Evidence that nerve growth factor dependence of sympathetic neurons for survival *in vitro* may be determined by levels of cytoplasmic free Ca. *Proc. natn. Acad. Sci. U.S.A.* **88**, 3892-3896.
- KOLB, B. & GIBB, R. (1991). Sparing of function after neonatal frontal lesions correlates with increased cortical dendritic branching: a possible mechanism for the Kennard effect. *Behav. Brain Res.* **43**, 51-56.
- KORN, H. & FABER, D. S. (1987). Regulation and significance of probabilistic release mechanisms at central synapses. In: *Synaptic Functions* (Edelman, W. M., Gall, W. & Cowan, W. M., eds) pp. 57-108. New York: John Wiley & Sons.
- LACAILLE, J.-C., MUELLER, A. L., KUNKEL, D. D. & SCHWARTZKROIN, P. A. (1987). Local circuit interactions between oriens/alveus interneurons and CA1 pyramidal cells in hippocampal slices: electrophysiology and morphology. *J. Neurosci.* **7**, 1979-1993.
- LANKDORF, K., DE MELLO, F. G. & KLEIN, W. L. (1987). A transient embryonic dopamine receptor inhibits growth cone motility and neurite outgrowth in a subset of avian retina neurons. *Neurosci. Lett.* **75**, 169-174.
- LIPTON, S. A. (1986). Blockade of electrical activity promotes the death of mammalian retinal ganglion cells in culture. *Proc. natn. Acad. Sci. U.S.A.* **83**, 9774-9778.
- LIPTON, S. A. & KATER, S. B. (1989). Neurotransmitter regulation of neuronal outgrowth, plasticity and survival. *Trends Neurosci.* **12**, 265-270.
- LU, B., YOKOYAMA, M., DREYFUS, C. F. & BLACK, I. B. (1991). Depolarizing stimuli regulate nerve growth factor gene expression in cultured hippocampal neurons. *Proc. natn. Acad. Sci. U.S.A.* **88**, 6289-6292.
- MADISON, D. V., MALENKA, R. C. & NICOLL, R. A. (1991). Mechanisms underlying long-term potentiation of synaptic transmission. *A. Rev. Neurosci.* **14**, 379-397.
- MATTSON, M. P. & KATER, S. B. (1989). Excitatory and inhibitory neurotransmitters in the generation and degeneration of hippocampal neuroarchitecture. *Brain Res.* **478**, 337-348.
- MCCAIG, C. D. (1990a). Nerve branching is induced and oriented by a small applied electric field. *J. Cell Sci.* **95**, 605-616.
- MCCAIG, C. D. (1990b). Nerve growth in a small applied electric field and the effects of pharmacological agents on rate and orientation. *J. Cell Sci.* **95**, 617-622.
- MCCOBB, D. P., HAYDON, P. G. & KATER, S. B. (1988). Dopamine and serotonin inhibition of neurite elongation of different identified neurons. *J. Neurosci. Res.* **19**, 19-26.
- MCDONALD, J. W. & JOHNSTON, M. V. (1990). Physiological and pathological roles of excitatory amino acids during central nervous system development. *Brain Res. Rev.* **15**, 41-70.
- MCDONALD, J. W., JOHNSTON, M. V. & YOUNG, A. B. (1990). Differential ontogenic development of three receptors comprising the NMDA receptor/channel complex in rat hippocampus. *Exp. Neurol.* **110**, 237-247.
- MILLER, M. W. (1988). Development of projection and local circuit neurons in neocortex. In: *Cerebral Cortex. Development and Maturation of the Cerebral Cortex*, Vol. 7. (Peters, A. & Jones, E. G., eds), pp. 133-175. New York: Plenum Press.
- MURRAY, J. D. (1989). *Mathematical Biology*. pp. 130-132. Berlin, Heidelberg: Springer-Verlag.
- O'BRIEN, R. J. & FISCHBACH, G. D. (1986). Excitatory synaptic transmission between interneurons and motoneurons in chick spinal cord cultures. *J. Neurosci.* **6**, 3284-3289.
- O'KUSKY, J. R. (1985). Synapse elimination in the developing visual cortex: a morphometric analysis in normal and dark-reared cats. *Dev. Brain Res.* **22**, 81-91.
- OPPENHEIM, R. W. (1991). Cell death during development of the nervous system. *A. Rev. Neurosci.* **14**, 453-501.
- PARNAVELAS, J. G., PAPADOPOULOS, G. C. & CANAVAGH, M. E. (1988). Changes in Neurotransmitters During Development. In: *Development and Maturation of Cerebral Cortex*, Vol. 7, *Cerebral Cortex* (Peters, A. & Jones, E. G., eds) pp. 177-209. New York: Plenum Press.
- PENTNEY, R. J. (1986). Quantitative analysis of dendritic networks of Purkinje neurons during aging. *Neurobiol. Aging* **7**, 241-248.
- PERETTO, P. & NIEZ, J.-J. (1986). Stochastic dynamics of neural networks. *IEEE Trans. Sys. Man. Cybernet.* **16**, 73-83.
- PRESS, W. H., FLANNERY, B. P., TEUKOLSKY, S. A. & VETTERLING, W. T. (1988). *Numerical Recipes in C, The Art of Scientific Computing*. Cambridge: Cambridge University Press.
- PRICE, D. J. & BLAKEMORE, C. (1985). Regressive events in the postnatal development of association projections in the visual cortex. *Nature, Lond.* **316**, 721-724.
- PURVES, D. & LICHTMAN, J. W. (1980). Elimination of synapses in the developing nervous system. *Science* **210**, 153-157.
- PURVES, D. (1988). *Body and Brain, A Trophic Theory of Neural Connections*. Cambridge, MA: Harvard.
- RAMAKERS, G. J. A., RAADSHEER, F. C., CORNER, M. A., RAMAKERS, F. C. S. & VAN LEEUWEN, F. W. (1991). Development of neurons and glial cells in cerebral cortex, cultured in the presence or absence of bioelectric activity: Morphological observations. *Eur. J. Neurosci.* **3**, 140-153.
- SCHILLING, K., DICKINSON, M. H., CONNOR, J. A. & MORGAN, J. I. (1991). Electrical activity in cerebellar cultures determines Purkinje cell dendritic growth patterns. *Neuron* **7**, 891-902.
- SCHREYER, D. J. & JONES, E. G. (1988). Axon elimination in the developing corticospinal tract of the rat. *Dev. Brain Res.* **38**, 103-119.
- SEGAL, M. S. & FURSHPAN, E. J. (1990). Epileptiform activity in microcultures containing small numbers of hippocampal neurons. *J. Neurophysiol.* **64**, 1390-1398.
- SHATZ, C. J. (1990). Impulse activity and the patterning of connections during CNS development. *Neuron* **5**, 745-756.
- SPITZER, N. C. (1991). A developmental handshake: neuronal control of ionic currents and their control of neuronal differentiation. *J. Neurobiol.* **22**, 659-763.
- STANFIELD, B. B. & O'LEARY, D. D. M. (1985). The transient corticospinal projection from the occipital cortex during the postnatal development of the rat. *J. comp. Neurol.* **238**, 236-248.
- STANTON, P. K. & SEJNOWSKI, T. J. (1989). Associative long-term depression in the hippocampus induced by hebbian covariance. *Nature, Lond.* **339**, 215-218.
- SUSSDORF, W. S. & CAMPENOT, R. B. (1986). Influence of the extracellular potassium environment on neurite growth in sensory neurons, spinal cord neurons and sympathetic neurons. *Dev. Brain Res.* **25**, 43-52.
- SWAAB, D. F., HOFMAN, M. A. & HONNEBIER, M. B. O. M. (1990). Development of vasopressin neurons in the human suprachiasmatic nucleus in relation to birth. *Dev. Brain Res.* **52**, 289-293.
- THOENEN, H. (1991). The changing scene of neurotrophic factors. *Trends Neurosci.* **14**, 165-170.
- TODD, R. D. (1992). Neural development is regulated by classical neurotransmitters: dopamine D2 receptor stimulation enhances neurite outgrowth. *Biol. Psychiatry* **31**, 794-807.
- TSUMOTO, T. (1992). Long-term potentiation and long-term depression in the neocortex. *Prog. Neurobiol.* **39**, 209-228.
- UYLINGS, H. B. M., VAN EDEN, C. G., PARNAVELAS, J. G. & KALSBECK, A. (1990). The prenatal and postnatal development of rat cerebral cortex. In: *The Cerebral Cortex of the Rat* (Kolb, B. & Tees, R. C., eds) Cambridge, MA: MIT Press.
- VAN HUIZEN, F. (1986). Significance of bioelectric activity for synaptic network formation. PhD Thesis, University of Amsterdam.
- VAN HUIZEN, F. & ROMIJN, H. J. (1987). Tetrodotoxin enhances initial neurite outgrowth from fetal rat cerebral cortex cells *in vitro*. *Brain Res.* **408**, 271-274.
- VAN HUIZEN, F., ROMIJN, H. J. & HABETS, A. M. M. C. (1985). Synaptogenesis in rat cerebral cortex is affected during chronic blockade of spontaneous bioelectric activity by tetrodotoxin. *Dev. Brain Res.* **19**, 67-80.



- VAN HUIZEN, F., ROMIJN, H. J., HABETS, A. M. M. C. & VAN DEN HOOFF, P. (1987). Accelerated neural network formation in rat cerebral cortex cultures chronically disinhibited with picrotoxin. *Exp. Neurol.* **97**, 280–288.
- VAN OOYEN, A., VAN PELT, J., CORNER, M. A. & LOPES DA SILVA, F. H. (1992). The emergence of long-lasting transients of activity in simple neural networks. *Biol. Cybern.* **67**, 269–277.
- VAUGHN, J. E. (1989). Review: fine structure of synaptogenesis in the vertebrate central nervous system. *Synapse* **3**, 255–285.
- VERVEEN, A. A. (1960). On the Fluctuations of Threshold of the Nerve Fibre. In: *Structure and Function of the Cerebral Cortex*, (Tower, D. B. & SCHADE, J. P., eds) Amsterdam: Elsevier.
- VON DER MALSBERG, C. & SINGER, W. (1988). Principles of Cortical Network Organization. In: *Neurobiology of Neocortex* (Rakic, P. & Singer, W., eds) pp. 69–99. John Wiley & Sons.
- WAHLE, P. & MEYER, G. (1987). Morphology and quantitative changes of transient NPY-ir neuronal populations during early postnatal development of the cat visual cortex. *J. comp. Neurol.* **261**, 165–192.
- WARREN, M. A. & BEDI, K. S. (1984). A quantitative assessment of the development of synapses and neurons in the visual cortex of control and undernourished rats. *J. comp. Neurol.* **227**, 104–108.
- WÄSSLE, L., PEICHL, L. & BOYCOTT, B. B. (1981). Dendritic territories of cat retinal ganglion cells. *Nature, Lond.* **292**, 344–345.
- WILSON, H. R. & COWAN, J. D. (1972). Excitatory and inhibitory interactions in localized populations of model neurons. *Biophys. J.* **12**, 1–24.
- ZAFRA, F., HENGERER, B., LEIBROCK, J., THEOEN, H. & LINDHOLM, D. (1990). Activity dependent regulation of BDNF and NGF mRNAs in the rat hippocampus is mediated by non-NMDA glutamate receptors. *EMBO J.* **9**, 3545–3550.

## APPENDIX

Derivation of eqn (8). Taking the average over  $i$  in eqn (7) and writing  $X_i = \bar{X} + \Delta X_i$ , with  $\bar{X} = (1/N) \sum_i^N X_i$ , yield

$$0 = -\bar{X}/\tau + \frac{1-\bar{X}}{N} \sum_j^N F(X_j) \sum_i^N W_{ij} - \frac{1}{N} \sum_j^N F(X_j) \sum_i^N \Delta X_i W_{ij}. \quad (\text{A.1})$$

Take  $\sum_i^N W_{ij} = N\bar{W}_{.j}$ , and write  $W_{ij} = \bar{W}_{.j} + \Delta_i W_{ij}$

$$0 = -\bar{X}/\tau + (1-\bar{X}) \sum_j^N F(X_j) \bar{W}_{.j} - \frac{1}{N} \sum_j^N F(X_j) \sum_i^N \Delta X_i \Delta_i W_{ij}. \quad (\text{A.2})$$

If  $\Delta X_i$  and  $\Delta_i W_{ij}$  are uncorrelated or small

$$0 \simeq -\bar{X}/\tau + (1-\bar{X}) \sum_j^N F(X_j) \bar{W}_{.j}. \quad (\text{A.3})$$

Take  $F(X_j) = F(\bar{X}) + \Delta X_j F'(\bar{X}) + \dots$ , and write  $\bar{W} = (1/N) \sum_j^N \bar{W}_{.j}$ ,  $\bar{W}_{.j} = \bar{W} + \Delta \bar{W}_{.j}$

$$0 \simeq -\bar{X}/\tau + (1-\bar{X}) F(\bar{X}) \bar{W} + (1-\bar{X}) F'(\bar{X}) \sum_j^N \Delta X_j \Delta \bar{W}_{.j} + \dots \quad (\text{A.4})$$

where  $\bar{W} = \sum_j^N \bar{W}_{.j} = (1/N) \sum_{ij}^N W_{ij}$ . If, again,  $\Delta X_j$  and  $\Delta \bar{W}_{.j}$  are uncorrelated or small, and ignoring higher order terms in the Taylor expansion

$$0 \simeq -\bar{X}/\tau + (1-\bar{X}) \bar{W} F(\bar{X}).$$